







- (1 60-69%):
10
- (1 50-
59%): 18
- (1 35-49%):
17
- (1
35%): 20

- 90,8%
(59 65);
- 56,9% (37) , 4 –
- 60% (39) 3 ;
- Candida
- 2




- S. Pneumoniae
- H. Influenzae
- M. Catarralis
- Enterobacteriaceae
- P. Aeruginosa
-

	(.)	(.)
S. Pneumoniae	25	10
H. influenzae	10	17
M. catarrhalis	4	6
Enterobacteriaceae	3	7
P. aeruginosa	2	6
	3	3





- S. Pneumoniae
- H. Influenzae
- M. Catarralis
- Enterobacteriaceae
- P. Aeruginosa
-



- S. Pneumoniae
- H. Influenzae
- M. Catarralis
- Enterobacteriaceae
- P. Aeruginosa
-


1

50%



- S. Pneumoniae
- H. Influenzae
- M. Catarralis
- Enterobacteriaceae
- P. Aeruginosa
-


1 35-50%



- S. Pneumoniae
- H. Influenzae
- M. Catarralis
- Enterobacteriaceae
- P. Aeruginosa
-

1

35%



■ S. Pneumoniae

■ H. Influenzae

■ M. Catarralis

■ Enterobacteriaceae

■ P. Aeruginosa



(%)

	1 > 50%	1 35- 50%	1 < 35%
	86	71	50
	85	72	61
	75	84	52
	98	90	75



.

;



.

;

-



,



<<

,

...>>